

The Phase Structure in the Evolution of Globally coupled Spatial Prisoner's Dilemma

Norihito Toyota and Shota Hayakawa

Hokkaido Information University, 59-2 Nishinopporo Ebetsu City
E-mail:toyota@do-johodai.ac.jp

Abstract

We propose two new evolutionary rules that is not mimic evolution of strategies based on the spatial Prisoner's Dilemma (PD). The former follows the selfish evolutionary rule and then the coexistence phase appears with weak phase transition with mutations. The second rule is that of globally coupled spatial PD where the strategy of an agent is taken so that the total payoff of the whole system increases. Then it is natulally suspected that all agents will become cooperators in general. We, however, find that situations is rather complex. We find two critical points within the usual conditions in the parameters included the payoff matrix in the PD. Further as we break the condition without losing PD condition, we meet two more essential curitical points. We discuss the areas between these critical points in the parameter space and uncover the various properties by simulating on computer. We show that in some parameter regions, strange properties, that is, like chaotic behaveor appear and the coexistence between defectors and cooperators appears in the large parameter region. Thus we point out that usual non-essential condition among the parameters in the payoff matrix imposed on the PD privent from recovering rich structures in the evolutionally spatial PS.

key words: *Prisoner's Dilemma, Evolutionary Game, Spational Game, Phase Structure, Globally Coupled, Critical Point*

1 Introduction

Traditionally differential equations have been applied to understand various phenomena in the nature and the artificial, especially physical, phenomena, since they were used to construct classical dynamics by I. Newton. It, however, seems that some phenomena are rather relevant to be studied as they follow some descrete dynamics. They may include biological, social, ecological phenomena and so on, which have a close relationship to "information". The game theory among many aproaches give one of interesting view of them. In this article we discuss the Prisoner's Dilemma game (PD), which is the most famous game and has substantial content among many sorts of games.

PD is symmetric game played by two agents and the payoff function acquired in each strategy is discribed by matrix form. One shot game of PD can be analyzed analytically [1]. Many researches have been made to resolve the Dilemma, *pareto optimal* $\neq$ *Nash equilibrium*, and how can we obtain cooperation rationally. Based on PD, iterated games have been studied and Axelrod [2] has shown that cooperation can emerge as a norm in a society comprised of individuals with selfish motives. Moreover spatial version of PG has been discussed so that a defection is to be only evolutionary stable strategy (ESS) [5] if each agent interacts with any other agent. This aspect drastically changes if a spatial structure of the population is considered. If the interaction between agents is locally restricted to their neighbors [3], a stable coexistence between cooperators and defectors become possible under certain conditions [3][4]. In the case, one agent plays PD with their neighbors and in next step the agent take the same strategy of the agent that acquired highest payoff among the neighbors, which reflects to Dawin's theory, sometimes "including mutation" [6]. Then it is assumed that

all agents play at same time and follow the same way. Recently the evolution of the spatio-structured PD has been systematically explored in details in Ref.[7].

We here discuss two other evolutionary rules within the framework of the spatio-structured PD where agents interact locally, in which the evolutionary rule entices the cooperator and defector to coexistence phase. One way to realize it is to change his(her) strategy when the agent would get more payoff if he(she) had changed the strategy. This selfish rule, however, leads to a trivial result, since PD has a dominant strategy. Second one is to change the strategy like totalitarian, that is, if the total payoff yielded in whole system increases when an agent changes the strategy. We call this evolutionary rule the globally coupled evolution. This evolutionary role is expected to lead to full cooperative strategy but it do not so. We found 4 critical points analytically, and perform computer simulations in the parameter regions bounded on the critical points.

After Introduction, section 1, we discuss our evolutionary strategy and the critical points analytically in the section 2. In the section 3 simulation results are obtained based on the results in the section 2. The final section 4 is devoted to summary.

2 Evolutionary Rule and Analytical Study

Spatially structured PD is first explained briefly. Then we discuss two evolutionary rules that are not considered thus far and study them analytically.

2.1 Spational Structured PD as Cellular Automaton

Two types of agents, cooperators (C) and defectors (D) are considered. The distinction is made by means of a variable $\theta \in \{-1, 1\}$, where 1 refers to cooperation and -1 refers to defection. An individual agent i is indicated by θ_i . N is total number of agents and the fraction of cooperators and defectors are given by

$$N = \sum_{\theta} N_{\theta} = N_{-1} + N_1 = \text{const}, \quad (1)$$

$$f_{\theta} = \frac{N_{\theta}}{N}; \quad f_{-1} + f_1 = 1. \quad (2)$$

The spatial distribution of the agents are considered as a two dimensional cellular automaton (CA) consisting of $N = n^2$ cells, where each cell is identified by the index $i \in N$ referring its spacial position and the state θ_i sits on the cell. The state space of all possible configurations is of the order 2^N .

We assume that each agent i has two options, C ($\theta = 1$) or D ($\theta = -1$). Playing with 4 neighbors of i , the action chosen by i and actions chosen by the 4 agents only affect the total payoff (Hamiltonian). We assume that an agent i simultaneously plays with 4 agents with a same strategy, and so the game essentially reduces to 2-person game. Then the two person game can be described by a payoff matrix and it is obtained by the following Table 1 for PD.

Table 1: Payoff table for PD. The left and right variables in the parenthesis show the payoffs of the agent i and j , respectively.

	C _j	D _j
C _i	(R, R)	(S, T)
D _i	(T, S)	(P, P)

In PD, the payoffs have to fulfill the following condition:

$$T > R > P > S, \quad (3)$$

and usually for iteration of game an additional condition is imposed:

$$2R > S + T. \quad (4)$$

The second condition, however, is considered to have not any essential meaning for the structured spatial game. Later the breakdown of this condition (4) induces some interesting phenomena in spatio PD. A simple analysis show that strategy D is so-called ESS in a one-shot PD [5]. It is also known that the pareto optimal strategy is for both agents to take C but Nash solution is for both agents to take D, which leads to famous "Dilemma".

In this paper we are interested in the spacial effects of PD and assume that each agent interacts only with the 4 agents of its neighborhood. In evolutionary game theory this is called a 4-person game. We here investigate the modified evolutionary rules, where players do not mimic the best strategy in its neighbors. We consider two extreme ways, one obeys a selfish evolutionary rule and another is like totalitarian, where explicit values of T, S, R, P play important role apart from the usual PD.

2.2 Selfish Evolution

Selfish evolution is to change strategy taken by an agent if the (target) agent get more payoff when he/she changes the strategy, independent of other agents. Suppose that all the agents follow this rule and each agent updates its strategy in regular order. We see this case leads to a trivial result.

In the simulation experiment of sec. 3 we include mutant agents with some probability with respect to Boltzmann distribution among regular agents that obey the selfish rule. This type agents will lead to the coexistence phase between C and D agents.

2.3 Globally Coupled Evolution

As the second evolutionary model, we consider that if the total payoff, hamiltonian, yielded in whole system increases when an agent changes the strategy, the agent assumes to change its strategy. So an agent decides its strategy according to the total payoff of the whole system. Due to this property, we call this model "globally coupled". Notice that the evolution of the system depends on from which agent the strategy is renewed at each round, since by changing a strategy of each agent the Hamiltonian generally is updated and the next target agent decides its strategy based on the updated Hamiltonian. It means that the order of agent targeted is important and the evolutionary process is only deterministic under a order. We mainly discuss the case of the following ordering;

$$[Do[Do[agent(i, j), \{i, 1, n\}], \{j, 1, n\}]]$$

in Mathematica notation, where (i, j) shows the coordinate (row element, column element) in lattice like CA. A random ordering will be also considered latterly.

2.4 Analytic study and critical points

We analytically investigate globally coupled evolution of PD in this subsection. Let consider the profit $P_i(C)$ that a cooperator agent i encircled by c cooperators and d defectors;

$$P_i(C) = cR + dS, \text{ with } c + d = 4. \quad (5)$$

If the agent change the strategy from C to D, the updated profit $P_i(D)$ of the agent is given by

$$P_i(D) = cT + dP, \quad (6)$$

and the difference is

$$\delta P(C \rightarrow D) = c(T - R) + d(P - S) > 0, \quad (7)$$

because of the equation (3). Noticing that the increment $\delta h(\theta_j)$ of the profit of a neighborhood j of the agent i brought by this change is

$$\delta h(\theta_j) = \begin{cases} S - R & \text{for } \theta_j = 1 \\ P - T & \text{for } \theta_j = -1, \end{cases} \quad (8)$$

the total $\delta h(\theta_j)$ resulted by 4 neighbors (c cooperators and d defectors) is given by

$$\Delta h = c(S - R) + d((P - T)) < 0. \quad (9)$$

This reverse of the strategy of agent i reflects the total profit (Hamiltonian) of the whole system and thus the resultant increment of the total Hamiltonian $\Delta H_{iC}[c, d]$ is given by

$$\Delta H_{iC}[c, d] = c(S + T - 2R) + d(2P - S - T), \quad (10)$$

where the first term in the right hand side is negative if the equation (4) is satisfied. In contrast with this when the agent i change the strategy from D to C, the corresponding amount becomes

$$\Delta H_{iD}[c, d] = -\Delta H_{iC}[c, d] \equiv -\Delta H[c, d] \quad (11)$$

by following the similar calculation. Explicitly they are written by

$$\Delta H[0, 4] = 4(2P - S - T), \quad (12)$$

$$\Delta H[1, 3] = 6P - 2(R + S + T), \quad (13)$$

$$\Delta H[2, 2] = 4(P - R) < 0 \text{ from (3)}, \quad (14)$$

$$\Delta H[3, 1] = 2(P + S + T - 3R) < 0 \text{ from (4)}, \quad (15)$$

$$\Delta H[4, 0] = 4(S + T - 2R) < 0 \text{ from (4)}, \quad (16)$$

where (14) is negative by definition of PD, and (15) and (16) are negative because of the additional condition (4). Whether (12) and (13) are positive or not depends on the explicit parameters even when satisfying (3) and (4). This point is different from usual analysis of PD where explicit values of P, R, S, T are not essential but the order of the magnitudes plays only important role. A standard set of parameters adopted by Okada [1] is $\{P, R, S, T\} = \{-3, 5, -4, 6\}$ and then all increments of the Hamiltonian turn are negative.

However, whenever the condition $2P < S + T$ is satisfied, $\Delta H[0, 4]$ and $\Delta H[1, 3]$ come to be positive, and moreover (15) and (16) become possible to be positive when the additional condition (4) is broken. To explain this situation, without losing the generality we can choose S and T as variable parameters while P and R are fixed to be the standard values. Namely as S decreases much, $\Delta H[0, 4]$ and $\Delta H[1, 3]$ increase to be positive and as T increases much, $\Delta H[3, 1]$ and $\Delta H[4, 0]$ increase to be positive. We find critical points in S when $\Delta H[0, 4]$ and (or) $\Delta H[1, 3]$ become 0, respectively. They turn out to be $S = -12$ and $S = -20$ whose values depend on the explicit values of P and R but this is not essential, because the fact that there are two critical points in any values of them and the signs of $\Delta H[0, 4]$ and $\Delta H[1, 3]$ change preserves to be true at these points. There are equally two critical points in T , when $\Delta H[3, 1]$ and $\Delta H[4, 0]$ cross zero point. They are $T = 14$ and $T = 22$. Thus the parameter space divides into five regions and they are summarized in the Fig.1.




Figure 1: Phase diagram

3 Simulation results

In this section the results of computer simulation on each region and the some discussions on them are obtained. We basically choose parameters according to Okada's book[1], i.e. $R = 5$, $P = -3$, $T = 6$, $S = -4$ and the lattice size $n = 13$ except for some special cases. The parameter S and T , however, are changed to be some correct values such as identifying the corresponding region. In the evolutionary PD, various types of initial configuration states are explored and many times experiments for a random initial state are made. The following initial configuration states are explored.

- (a) Initial configuration has only defectors.
- (b) Initial configuration has only defectors except one cooperator.
- (c) Initial configuration has only cooperators except one defector.
- (d) Initial configuration has only cooperators.
- (e) Initial configuration has cooperators and defectors at random positions where the ratio $\frac{N_1}{N-1}$ of populations between cooperators and defectors is parametrized by $p(0 \leq p \leq 1)$.

Though the lattice size $n = 13$ seems to be too small, we have ascertained that essential results are unchanged by magnifying the lattice size except for some cases pointed specially. While states immediately converge in almost every case, some interesting cases show rather complex behaviors such as chaotic-like behaviors.

3.1 Selfish Evolution

As all the agents follow this rule, full agents become defectors in the second step (i.e. after one round) because the dominant strategy in PD is a defection. Simulation results of course supports this argument.

This situation is unchanged even in introducing a stochastic process by the Metropolis method with Boltzmann distribution, where player i evade the above evolutionary rule with the probability $1 - e^{-q\delta P_i}$, at low temperature. (q is inverse of temperature.) At high temperature, however, a coexistence phase between cooperators and defectors appears. The critical value is $q = 1.0 \sim 2.0$, that is, when $q > 2$ the system is covered to defectors after a few steps. Between $q = 1.0$ and $q = 2.0$ the system is unstable, and all defectors age and the age with a few cooperators revolve. As q decrease from 1.0 to 0, the average of the population of cooperators slowly increases to the half of total population with some fluctuation. A typical example with $q = 0.2$ is given in the Fig. 2.

These results also do not depend on the initial state of the system, starting even from full defection (cooperation) initially.

3.2 Globally Coupled Evolution

(i)Region A

$\Delta H[4, 0] > 0$, $\Delta H[3, 1] > 0$ in this region. Thus an agent surrounded by many (3 and 4) C-agents becomes D-agent but one surrounded by small number of C-agents remains C-agent at the next round. The coexistence of C and D as the final state is predicted. We simulate the case with $T = 25$ and $T = 50$. It turns out that independently of an initial state, the population ratio between C and D is $N_1 : N_{-1} = 1 : 1$ within 3% error (for random initial state, within 10% error) after a few rounds. A typical example is given in the Fig. 3.

(ii)On the boundary between A and B

$\Delta H[4, 0] > 0$, $\Delta H[3, 1] = 0$ in this region with $T = 22$. Agents take the same action as the region A, since we assume that an agent also change its strategy when $\Delta H = 0$. Taking (a) and (b) initially, the state results in $N_1 : N_{-1} = (a + n) : a = (n + 1) : (n - 1)$ with $n + 2a = n^2$ by a simple analytic calculation from the resultant diagram after about n rounds (Fig.4). Thus as $n \rightarrow \infty$, N_1 asymptotically comes close to N_{-1} .

However this convergence is so unstable that the state does not converge even when initial state has two C-agents. Interestingly N_1 is exactly periodic and during the periodic age the total Hamiltonian invariant, while the states itself is changing.

Random initial states with $p = 0.5$ also expose almost convergent behaviour in N_1 but they are exactly periodic really (Fig.5). The period is scale dependent.

Generally various types of periodic behaviours in N_1 appear in random initial configurations but the configurations of the state never converge and so are not periodic (of course strictly speaking, it is periodic because the dimension of the configuration space in size n is 2^N).

For (d) and (c), N_C shows an intermittent but exactly periodic behaviour. The period is scale dependent and $n + c$ with the uncertain origin of the constant $c \sim 14$ (fig.6).

(iii)Region B

In this region $\Delta H[4, 0] > 0$ and so an agent surrounded by 4 C-agents only converts to D-agent. We choose $T = 16$ as a typical value. For $n = 13$ the state rapidly converge to be $N_C/N_D = 1.63$. By analytical consideration we easily find that $\frac{N_1}{N_{-1}} = \frac{(n^2+2n-3)}{(n^2-2n+3)} \equiv r$ with $N_1 - N_{-1} = 2n - 3$ and thus its ratio approaches to 1 as $n \rightarrow \infty$. This result is common in the cases, (a),(b),(c),(d) as in the Fig.7, while the ratio fluctuates in the cases with random initial configurations. Although the state at $0 \leq p < 0.2$ and $0.8 < p \leq 1.0$ converges to be the ratios $\sim r$, in the cases with $0.2 \leq p \leq 0.8$ the ratios are enhanced to the ratios with $N_1 - N_{-1} \sim 4n$.

(iv)On the boundary between B and C

This region is identified by $\Delta H[4, 0] = 0$ with $T = 14$ and all other $\Delta H < 0$. Except for the cases with the random initial configurations, N_1 vibrates with round-dependent amplitude. Thus the amplitude itself vibrates with some long period (Fig.8), while the configurations show highly geometrical patterns.

For $n = 13$ the period is 23 ± 1 (essentially $2(n - 1)$) when states initially begin with (a),(b),(c),(d) and the total Hamiltonian remains invariant during the period. On the other hand, beginning with random initial state, the evolution of the state is prima facie chaotic (Fig.9). However, constructing the return map, we can find a weak correlation in N_1 data (the Fig.10). It is conjectured that the initial randomness only weakly breaks the above periodic property.

(v) Region C (standard region)

In this case we have all $\Delta H < 0$. As a typical value of S and T , $S = -4$ and $T = 6$ are taken. As expected, the state rapidly converges to the state with all C, independent of initial configuration states.

(vi) On the boundary between D and C

Except for $\Delta H[0, 4] = 0$, all ΔH is negative. In this case the state rapidly converges to the state with all C with little marvel. This depends on the ordering of target agent as above mentioned. If C agent surrounded by 4 D-agents is lackily a first target in D dominant worlds, this situation will change drastically (this can be also confirmed by computer simulation really). In this simulation it is not so, that is, an agent D next to C agent first is remarked to turn to C agent and so the original C agent can not become D agent when it is remarked. Thus C agents only can live in the final state. This arises except for the case (a) where the state with all R is stable.

(vii) Region D

In this region with $S = -18$ we have only positive $\Delta H[0, 4] > 0$. In this time an agent around without any C agent only becomes D agent. With the same reason as (vi) essentially, C agents only can live in the final state, while the state (a) with all D agents anomalously is stable.

(viii) On the boundary between D and E

In this case we take $S = -22$, and $\Delta H[1, 3] = 0$ and $\Delta H[0, 4] > 0$. The phenomenological results are exactly same as the cases of (vii).

(ix) Region E

In this region with $S = -25$ we have two positive $\Delta H[0, 4] > 0$ and $\Delta H[1, 3] > 0$. The phenomenological results are exactly same as the case of (vii) except one problematic point. In the case of (b), the state drops into the state with all D agents. Since the random initial state with $p = 0.5$ finally comes to the state with all D agents, when the number of initial C agents decreases, there should be some kind of phase transition. We study this by varying the initial ratio f_1 , that is to say, the parameter p . They are summarized in the figure 11. Around $p = 0.2$, we find that a weak phase transition occurs.

3.3 Some Remarks

In the selfish evolution model without mutants, the state trivially comes to be full D-agent dominated world. In the case with mutants, however, there is a coexistence phase between D and C. As varying the mutation ratio

q , a sort of phase transition appears at around $q = 1.5 \pm 0.5$. When $q < 1.0$, it is inferred that the population ratio N_1/N_{-1} is correlated with q , that is, the degree of randomness, from the simulation results, and thus the results may be almost consequential one.

In the globally coupled model, any interesting phenomena do not arise at critical points related to $\Delta H[1, 3]$ and $\Delta H[0, 4]$, while rich structures appear at critical points related to $\Delta H[3, 1]$ and $\Delta H[4, 0]$ which emerges at the regions breaking the condition (4). There are the coexistence phase in the latter regions, widely.

This can be understood by considering the reduced payoff table which is constructed by adding the payoffs of two agents at each element of the payoff table 1. This can be also derived by noticing that S and T always appears pairwise and P and R always appears as multiples of 2 in ΔH 's. The reduced table for PD is obtained in the Table 2. Basically agents tend to be a C-agent in the globally coupled rule (that is, (C,C) is favored) with the total payoff $2R$ from the table 2. This tendency is broken at $2R < T + S$ and the agents' preference is changed to the strategy D from C. This consideration give the two effective critical points related to $\Delta H[3, 1]$ and $\Delta H[4, 0]$. On the other hand, $S + T < 2P$ is irrelevant to the change of the tendency. Thus the critical points related to $\Delta H[1, 3]$ and $\Delta H[0, 4]$ do not contribute to this story drastically.

Table 2: Reduced payoff table for PD.

	C_j	D_j
C_i	$(R + R)$	$(S + T)$
D_i	$(T + S)$	$(P + P)$

To eliminate the order dependence of a target agent, we choose target agent at random to perform the same simulation. It, however, turns that the results are unaffected by the order essentially. The longer period is merely needed untill the state converges. But, almost chaotic behaviour that does not show any regular behavior in the 2 and 3 dimensional embedding space appears in the region B with random initial state, which should be analyzed in details.

4 Summary

In this article we propose two new evolutionary rules in which the explicit values in the payoff table play important roles apart from usual games. By these we mainly investigate the evolution of the spatio PD.

In the selfish evolutionary rule with mutations, coexistence phase appears with weak phase transition where the order parameter is p . This may be almost understood trivially.

In the globally coupled rule it is natural to expect that a full D agent world is realized. As not expected, the rule presents rich structures in the spatio evolutionary PD. Analytically we find four critical points. When the condition $S + T < 2R$ usually imposed is broken that is related two critical points, the coexistence phase appears in the wide range of parameter space, while the other two critical points do not any special role in the evolution (as expected all C world is generally generated), except for region E. The region E shows a little strange feature where there is a transition between all C world and all D world.

Thus we uncover the entire phase structures under these rules. Detail analysis for some complex data in N_1 such as chaotic behavior remain to us.

References

- [1] A. Okada, "Game Theory", Yuuhikaku 1996 (in Japanese)

- [2] R. Axelrod, "The evolution of Cooperation", Basic Books, New York, 1984
- [3] M.A.Nowak and R. M. May, "The spatial dilemmas of evolution", Int. Jour. of Bifurcation and Chaos, 3(1), pp.35-78, 1993
- [4] M.A.Nowak and R. M. May, "Evolutionary games and spatial chaos", Nature, 359,pp.826-829, Oct. 1992
- [5] J. M. Smith, "Evolution and the theory of games" Cambridge University Press, UK, 1982
- [6] H. Fort, "Exploring the cooperative regimes in a model of agents without memory or "tag": indirected reciprocity vs. selfish incentives", available from <http://xxx.lang.gov/abs/nlin.AO/0211024>
- [7] F. Schwetzer, L. Behera and H. Muhlenbein, "Evolution of Cooperation in a Spatial Prisoner's Dilemma", available from <http://xxx.lang.gov/abs/cond-mat/0211605>




Figure 2: Coexistence state diagram of C and D with $q=0.2$ where black lattices denote D-agents and white ones do C-agents.




Figure 3: Coexistence state diagram in the region A with initial (a) and $T = 25$.




Figure 4: Coexistence state diagram on the boundary between B and C with initial (c) and $T = 22$




Figure 5: The number of C-agents on the boundary between B and C with $q=0.5$ random initial state and $T = 22$




Figure 6: The number of C-agents on the boundary between B and C with initial (a) and $T = 22$.




Figure 7: Coexistence state diagram in the region B with $T = 16$ and $n = 51$




Figure 8: N_C sequence on the boundary between B and C with initial (a) and $T = 14$




Figure 9: N_C sequence on the boundary between B and C in the random initial state with $p = 0.5$ and $T = 14$




Figure 10: Return Map on the region between B and C in the random initial state with $p = 0.5$ and $T = 14$




Figure 11: q vs. N_C and N_D diagram in the random initial state with $S = -25$